

Remission of Proteinuria Improves Prognosis in IgA Nephropathy

Heather N. Reich,* Stéphan Troyanov,[†] James W. Scholey,* and Daniel C. Cattran,* for the Toronto Glomerulonephritis Registry

*Division of Nephrology, University Health Network, University of Toronto, Toronto, Ontario, and [†]Department of Medicine, Division of Nephrology, Hôpital du Sacré-Coeur de Montréal, Faculty of Medicine, Université de Montréal, Montreal, Quebec, Canada

ABSTRACT

Proteinuria has been shown to be an adverse prognostic factor in IgA nephropathy. The benefit of achieving a partial remission of proteinuria, however, has not been well described. We studied 542 patients with biopsy-proven primary IgA nephropathy in the Toronto Glomerulonephritis Registry and found that glomerular filtration rate (GFR) declined at -0.38 ± 0.61 ml/min per $1.73 \text{ m}^2/\text{mo}$ overall, with 30% of subjects reaching end-stage renal disease. Multivariate analysis revealed that proteinuria during follow-up was the most important predictor of the rate of GFR decline. Among the 171 patients with <1 g/d of sustained proteinuria, the rate of decline was 90% slower than the mean rate. The rate of decline increased with the amount of proteinuria, such that those with sustained proteinuria >3 g/d ($n = 121$) lost renal function 25-fold faster than those with <1 g/d. Patients who presented with ≥ 3 g/d who achieved a partial remission (<1 g/d) had a similar course to patients who had ≤ 1 g/d throughout, and fared far better than patients who never achieved remission. These results underscore the relationship between proteinuria and prognosis in IgA nephropathy and establish the importance of remission.

J Am Soc Nephrol 18: 3177–3183, 2007. doi: 10.1681/ASN.2007050526

Primary IgA nephropathy (IgAN) is the most common form of idiopathic glomerulonephritis (GN) throughout the world and the main cause of ESRD in patients with primary glomerular disease.¹ In Toronto, nearly 40% of patients with IgAN progress to ESRD by 10 yr.² Variability in the literature regarding the clinical course of patients with IgAN^{3–7} may be related to multiple factors, including patient biologic differences, nephrologists' practice patterns (e.g., biopsy timing), and geography.⁷ The ability to predict outcome in patients with IgAN remains a critical feature of patient treatment and has been a primary focus of previous studies.^{2,3,5,6,8–15}

Many studies^{7,16,17} have shown that proteinuria is a predictor of outcome in IgAN. Experimental evidence supports these human data by clarifying the direct deleterious effect of proteinuria on renal tissue.^{18–23} In contrast to other progressive types of GN, in which it seems that only sustained nephrotic-range proteinuria (>3 to 3.5 g/d) ensures a poor prognosis, studies

in IgAN^{5,10,17,24,25} and have suggested that much lower levels of proteinuria adversely affect prognosis. Furthermore, although the concept of achieving "partial remission" is not commonly associated with IgAN, recent evidence in other forms of GN has determined its clinical significance and established its prognostic value.^{26,27} The relevance of such a definition and its value in IgAN would be important for management of this disease.

Accordingly, we examined the effects of proteinuria at diagnosis as well as sustained exposure to

Received May 1, 2007. Accepted July 4, 2007.

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Dr. Heather Reich, c/o The Toronto Glomerulonephritis Registry, Toronto General Hospital, 12EN-228, 200 Elizabeth Street, Toronto, Ontario M5G 2C4, Canada. Phone: 416-340-4187; Fax: 416-340-3714; E-mail: h.reich@utoronto.ca

Copyright © 2007 by the American Society of Nephrology

proteinuria on outcome in a large cohort of patients who had primary IgAN and were followed longitudinally in the Toronto Glomerulonephritis Registry. We also assessed the prognostic relevance of achieving partial remission as defined by achieving sustained proteinuria values <1 g/d.

RESULTS

Baseline Characteristics and Clinical Outcome

As described in Table 1, the majority (61%) of the 542 patients studied were male and of white or Asian (China, Japan, or Pacific Rim) descent. Mean initial proteinuria was 2.4 g/d, and creatinine clearance (CrCl) was 77 ml/min per 1.73 m². Patients were followed for 6.5 yr. By study completion, one third of patients had reached ESRD. The rate of progression as measured by slope of CrCl was -0.38 ± 0.61 ml/min per 1.73 m²/mo (-4.56 ml/min per 1.73 m²/yr).

Proteinuria and the Rate of Decline of Renal Function

Clinical variables at onset of disease and during follow-up were tested for association with the slope of CrCl. Regression anal-

ysis revealed that several factors, shown in Table 2, were predictive of a faster rate of renal function decline (steeper, more negative slope).

The time-average proteinuria (TA-proteinuria; see Concise Methods section) was a critical determinant of slope by univariate and multivariate analysis ($P < 0.01$) and the most important predictor of renal function decline ($R^2 = 0.162$, $F = 104.5$, $P < 0.01$). Proteinuria at presentation was not predictive of slope by multivariate analysis.

The TA-proteinuria was also analyzed as a categorical variable, as illustrated in Table 3. When adjusted for multiple comparisons, the rate of deterioration (slope) of renal function differed significantly across the entire range of proteinuria. The greatest difference in rate of decline occurred between the ≤ 1 and > 1 g/d TA-proteinuria. On the basis of the mean 10-yr slope, patients with ≤ 1 g of TA-proteinuria had stable renal function, and patients with 1 to 2 g of TA-proteinuria had a projected unadjusted loss of 40 ml/min per 1.73 m² by 10 yr. Trend test and nonparametric testing did not establish differences among patients with ≤ 1 g/d TA-proteinuria: 0 to 0.3 ($n = 36$) versus 0.3 to 0.6 ($n = 63$) versus 0.6 to 1 g/d TA-proteinuria ($n = 72$; slopes 0.00 ± 0.39 , 0.02 ± 0.48 , and -0.06 ± 0.48 ml/min per 1.73 m²/mo, respectively; NS). Proteinuria categories did not overlap (*i.e.*, patients were assigned to only one category of TA-proteinuria).

Table 1. Baseline characteristics and clinical outcome in 542 patients with IgAN

Characteristic	Value
Baseline	
age (yr; mean \pm SD [range])	38.2 \pm 13.01 (15.6 to 73.5)
gender (% male)	61
BMI (kg/m ² ; mean \pm SD [range])	25.9 \pm 5 (16 to 43)
creatinine (μ mol/L; mean \pm SD [range])	129.7 \pm 70.6 (50 to 731)
MAP (mmHg; mean \pm SD [range])	103 \pm 16 (73 to 170)
24-h urine protein (g; mean \pm SD [range])	2.37 \pm 2.5 (0 to 22)
CrCl (ml/min per 1.73 m ² ; mean \pm SD [range])	77.0 \pm 33 (9 to 217)
ethnicity (%)	
white	50.4
black	2.8
Asian	23.1
other	10.0
unknown	13.7
Follow-up	
length of follow-up (mo; mean \pm SD [range])	78.1 \pm 59 (12 to 315)
MAP (mmHg; mean \pm SD [range])	99.7 \pm 10 (72 to 141)
peak proteinuria (g/24 h; mean \pm SD)	3.71 \pm 3.20
proteinuria (g/24 h; mean \pm SD)	2.19 \pm 1.94
treatment (%)	
ACEi or ARB	53
fish oil	15.1
any immunosuppression	15.7
prednisone	12.5
renal failure (%)	27.7
slope CrCl (ml/min per 1.73 m ² /mo; mean \pm SD)	-0.38 ± 0.61

Proteinuria and Renal Survival

The TA-proteinuria was the most important predictor of renal survival, even when corrected for other parameters (multivariate hazard 1.57; 95% confidence interval 1.39 to 1.77; $P < 0.01$). The graded effect of TA-proteinuria on renal survival is illustrated in Figure 1. Having < 0.3 g/d proteinuria was similar to 0.3 to 1.0 g/d, but each gram above 1 g/d (reference group) was associated with worse renal survival, with a hazard ratio of 3.5 for ESRD with 1 to 2 g/d, 5 with 2 to 3 g/d, and 10 with > 3 g/d.

We analyzed the course of patients whose initial proteinuria was > 1 g/d and fell below this level during follow-up. Patients were divided into groups on the basis of their peak proteinuria: 1 to 2, 2 to 3, and > 3 g/d. As illustrated in Figure 2, all patients whose proteinuria reached a “partial remission” of < 1 g/d, regardless of the starting point, had a similar, favorable survival (log rank NS) and rate of progression (-0.2 ± 0.47 , -0.14 ± 0.63 , and -0.16 ± 0.37 ml/min per 1.73 m²/mo respectively; NS). In comparison, the outcome of “nonremitters” (*i.e.*, those who did not achieve < 1 g/d) was markedly worse than remitters, with a far more rapid rate of renal decline (-0.76 ± 0.59 versus -0.13 ± 0.41 ml/min per 1.73 m²/mo respectively; $P < 0.01$). Patients who achieved complete remission of proteinuria (sustained level of < 0.3 g/d with preserved CrCl) did not have a different rate of renal function decline than patients who achieved partial remission to < 1 g/d (-0.17 ± 0.47 versus -0.13 ± 0.41 ml/min per 1.73 m²/mo, respectively; NS).

The course of patients whose initial proteinuria was < 1 g/d but subsequently rose was also analyzed. Change in proteinuria was calculated by subtracting the final from the initial value. Patients

Table 2. Factors at presentation and during follow-up influencing decline in renal function (slope in ml/min per 1.73 m²/mo) by univariate and multivariate regression^a

Parameter	Univariate		Multivariate	
	Mean/Standardized β	P	Mean/Standardized β	P
Presentation				
ln(24 h Upro) ^b	-0.145/-0.258	<0.01	-	NS
MAP	-0.005/-0.145	<0.01	-	NS
Follow-up				
ln(24 h Upro) ^b	-0.258/-0.403	<0.01	-0.302/-0.493	<0.01
MAP	-0.020/-0.337	<0.01	-0.013/-0.231	<0.01
BP med	-0.088/-0.105	0.02	-	NS
ACEi/ARB ^c	0.227/0.092	0.03	0.077/0.124	0.02

^aUpro, urine protein excretion; BP med, average number of BP medications during follow-up.

^bUrine protein results were log transformed.

^cACEi/ARB exposure was measured as a continuous TA value and analyzed in quartiles of medication exposure (see Concise Methods section).

Table 3. Outcome based on categorical grouping of TA 24-h urine protein excretion during follow-up^a

Group (n)	TA-Proteinuria (g/24 h)	Slope (ml/min per 1.73 m ² /mo; Mean \pm SD)	Renal Failure Risk (Hazard [95% CI])
1 (171)	0 to 1	-0.030 \pm 0.46	Reference
2 (145)	1 to 2	-0.326 \pm 0.53	3.48 (1.8 to 6.7)
3 (105)	2 to 3	-0.516 \pm 0.66	5.17 (2.6 to 10.0)
4 (121)	>3	-0.719 \pm 0.61	9.89 (5.3 to 18.4)

^aCI, confidence interval.

were then grouped into three categories according to their quantitative increase in proteinuria: 1 to 2 g/d (group 1), 2 to 3 g/d (group 2), and >3 g/d (group 3). As illustrated in Figure 3, the opposite of the improvements seen with lowering proteinuria were observed: The greater the rise in proteinuria, the worse the renal survival (log rank $P = 0.004$ by trend test) and rate of renal function decline ($F = 3.8$, $P = 0.025$). *Post hoc* analysis revealed that although the rate of renal function decline of patients in group 1 was similar to that of group 2 (-0.23 ± 0.41 versus -0.12 ± 0.79 ml/min per 1.73 m²/mo; NS), patients in group 3 had a significantly more rapid rate of renal function decline than patients in either group 2 (-0.51 ± 0.55 versus -0.12 ± 0.79 ml/min per 1.73 m²/mo; $P = 0.02$) or group 1 (-0.51 ± 0.55 versus -0.23 ± 0.41 ml/min per 1.73 m²/mo; $P = 0.03$). In terms of renal survival, 20 of the patients who presented with “low risk” proteinuria levels of <1 g/d had a subsequent rise in proteinuria and actually progressed to ESRD. These 20 patients had a lower CrCl at presentation (58 versus 81.4 ml/min per 1.73 m²; $P < 0.05$) and a higher TA mean arterial pressure (TA-MAP; 101 versus 96 mmHg; $P < 0.05$).

Role of Angiotensin-Converting Enzyme Inhibitor/Angiotensin Receptor Blocker

Because this is not a therapeutic trial, the direct role of specific interventions cannot be accurately assessed. It was observed that the use of angiotensin-converting enzyme inhibitor/angiotensin receptor blocker (ACEi/ARB) lowered the rate of decline of renal function, even when adjusted for other clinical parameters. A greater proportion of patients who achieved partial remission of proteinuria were treated with ACEi/ARB, compared with patients who did not (66 versus 52%; $\chi^2 = 12.7$, $P < 0.01$). The beneficial effects of ACEi/ARB may have been

related to effects on both proteinuria and BP; however, use of other antihypertensive agents did not influence loss of renal function by multivariate analysis. The use of ACEi/ARB and GFR at the start of these medications were significant determinants of renal survival by univariate analysis ($P < 0.01$), however not multivariate analysis. The majority (86%) receiving ACEi/ARB were treated with ACEi alone, and there were insufficient patient numbers to analyze relative differences in ACEi versus ARB versus combination therapy.

Other Determinants of Outcome

Multivariate analysis revealed that only TA-proteinuria, TA-MAP, and quartile of exposure to ACEi/ARB were predictors of renal function decline (see Table 2) and were also predictive of






Figure 1. Renal survival by category of TA-proteinuria.



Group 1	29	18	6
Group 2	21	13	2
Group 3	39	26	10

Figure 2. Impact of reduction of proteinuria on renal survival. Patients attaining partial remission (<1 g/d), regardless of peak proteinuria, had a similar favorable outcome (NS). Group 1, 1 to 2 g/d peak proteinuria; group 2, 2 to 3 g/d peak proteinuria; group 3, >3 g/d peak proteinuria.

renal survival by multivariate Cox regression analysis. The complex nature of the interaction between MAP and proteinuria is illustrated in Figure 4; patients with the highest quartile of TA-MAP also had high levels of proteinuria and the greatest rate of loss of renal function; however, TA-proteinuria was still the most important predictor of outcome, independent of BP. Neither body mass index (BMI; continuous variable) nor high BMI (>27) was a predictor of slope by univariate analysis. As a continuous



Group 1	37	27	11
Group 2	16	8	2
Group 3	37	25	11

Figure 3. Renal survival in patients with ≤ 1 g/d proteinuria at presentation differed significantly on the basis of increase in proteinuria from baseline. Group 1, 1 to 2 g/d increase; group 2, 2 to 3 g/d increase; group 3, >3 g/d increase in proteinuria from baseline. $P = 0.004$ by trend test.




Figure 4. Relationship between TA-proteinuria group, quartile of MAP, and loss of CrCl. Group 1, TA-proteinuria <1 g/24 h; group 2, 1 to 2 g/24 h; group 3, 2 to 3 g/24 h; group 4, >3 g/24 h.

variable, BMI related to renal survival only by univariate (not multivariate) Cox regression, which may have been driven by a higher risk for ESRD in patients with BMI >29 (hazard ratio 2.1; 95% confidence interval 1.2 to 3.6). No other factor (initial renal function, smoking, gender, ethnicity, fish oil, or immunotherapy) was predictive of the rate of loss of kidney function.

DISCUSSION

The purpose of this study was to quantify the value of proteinuria reduction on outcome in patients with IgAN. We also sought to determine whether by defining partial remission and by assessing the impact of change in proteinuria on outcome we could confirm the importance of the definition by quantifying its value in relationship to both rate of disease progression and renal survival.

Although proteinuria is a known risk factor for progression of IgAN,^{2-6,8,9,14,28} important questions regarding its role in the prognosis of IgAN remain. First, timing of measurement requires clarification; proteinuria at diagnosis often is not a predictor of outcome by multivariate analysis,^{2,17} whereas proteinuria at 1 yr or later may better indicate prognosis.² Second, although many studies suggest that patients with ≤ 1 g/d at presentation have a favorable prognosis,^{5,29} this observation is not uniform,¹⁰ and it is not known whether patients who achieve this target from higher values have the same prognosis as patients who present with and maintain low-level proteinuria. Finally, the importance of defining partial remission in proteinuria was established recently for other forms of GN^{26,27}; it has not been confirmed or quantified in IgAN and would represent important information for clinicians.

We have confirmed that in IgAN, proteinuria exposure over time (TA-proteinuria) is the strongest predictor of the rate of renal function decline. The relationship between TA-proteinuria and outcome is dramatically altered down to levels as low as 1 g/d, which is in marked contrast to the other progressive types of primary nephropathy, including membranous GN and FSGS. A quantitative estimate of the impact of proteinuria

has been determined: Each incremental gram per day above 1 is associated with a 10- to 25-fold more rapid rate of renal function decline and similar differences in renal survival. Although we could not determine a difference in outcome below 1 g/d, this may reflect inadequate statistical power because the benefit is likely continuous; however, even if power were improved, the clinical relevance of improved slope below this level will be minor given the overall slow rate of decline observed in the patients with <1 g/d TA-proteinuria.

Patients who reached <1 g/d proteinuria regardless of their starting point, whether “partial remission” was reached spontaneously or with intervention, had an excellent prognosis, similar to patients whose proteinuria never exceeded 1 g/d. This is strong support for using this partial remission definition as a goal for clinicians and provides new insights regarding the value of proteinuria reduction as a therapeutic target. Although previously recognized as important, the magnitude of the effect of proteinuria reduction on progression and renal survival has not been described in IgAN. By determining the quantitative value of partial remission, intensive therapy targeting significantly lower levels of proteinuria than in the other primary glomerulopathies is justified. Providing this information to the patient—that is, the substantial value of small reductions in proteinuria—should also aid compliance in these largely asymptomatic individuals.

This study was not a therapeutic trial and not designed to assess the benefit of therapeutic interventions in the course of IgAN. The most important conclusion derived is that the reduction of proteinuria by whatever means (medication, MAP reduction) is of great clinical benefit. The use of ACEi/ARB was the only intervention associated with a slower rate of renal function decline and prolonged renal survival, in keeping with both published randomized, controlled trials and retrospective reviews demonstrating the renoprotective effect of renin-angiotensin-aldosterone system blockade.^{30–39} These beneficial effects may reflect both hemodynamic and nonhemodynamic antiproteinuric effects of ACEi/ARB^{24,35,40–42} or alterations in glomerular permselectivity⁴³; however, this study was not designed to determine causality or mechanism. Similarly, although MAP was an important determinant of outcome and patients with the highest quartile of MAP had the highest levels of proteinuria, this is not necessarily a causal relationship.

The results of this study may not be generalizable to other populations. We previously demonstrated differences across geographic regions of the world in patients with IgAN.⁷ Although the majority of the difference was due to practice patterns, we could not explain all of the regional differences. In addition, because of the dominant effect of protein reduction on outcome, the effects of interventions and clinical parameters (fish oil, BMI) may have been obscured. Finally, this study was not designed to assess the role of renal biopsy findings in predicting outcome. Our aim was to measure factors during the course of disease that relate to outcome (e.g., TA-proteinuria), as opposed to implications of cross-sectional factors, such as biopsy grade at the time of diagnosis. As demonstrated in our previous study² of a large cohort of patients

with IgAN, although biopsy findings are clinically important, we do not believe that the findings at the time of biopsy provide additional information that is not captured by time-dependent variables. Although many studies have found a relationship between sclerosis/fibrosis and outcome, these studies did not include sequential measurements of MAP/proteinuria over time^{5,44,45} or had shorter follow-up.³ Indeed, a relationship between biopsy class and final follow-up data has been noted¹⁵; however, the value of adding the pathologic information to repeated sequential clinical measurements over time has not been demonstrated.

In summary, sustained proteinuria >1 g/d was the strongest predictor of the rate of progression of renal disease and the development of renal failure in IgAN. We demonstrated that with each sustained gram-per-day increment of proteinuria above 1, fold differences in progression rate and renal survival were observed. More important, patients who were able to achieve and sustain reduction in proteinuria to <1 g/d had an excellent prognosis regardless of the level of initial proteinuria. This study quantifies the impact of proteinuria reduction in IgAN and the clinical relevance of defining partial remission in this disease as a valuable prognostic indicator for both the clinician and the patient.

CONCISE METHODS

Patient Selection

As described previously,^{26,27,46} the Toronto Glomerulonephritis Registry was started in 1974 and includes all biopsy-proven cases of GN from the greater Toronto area. Patient information is documented from first clinical presentation and collected on a periodic prospective basis by registrars.

All patients who had biopsy-proven IgAN and were enrolled in the Toronto Glomerulonephritis Registry were considered ($n = 1373$) and were excluded only when clinical data were incomplete (37 lacked proteinuria data, 18 lacked weight data), they were younger than 16 yr at presentation ($n = 54$), they had <12 mo of follow-up ($n = 713$), or they had a secondary cause of IgA deposition ($n = 9$). A total of 542 patients were included.

Gender, ethnicity, age, and BMI were recorded at the time of first assessment suggestive of GN. Weight; BP; exposure to medications; and laboratory parameters recorded, including creatinine, albumin, urinalysis results, and 24-h urine protein and creatinine excretion, were collected prospectively.

Definitions

CrCl was estimated using the Cockcroft-Gault method adjusted for body surface area.^{47,48} The GFR was also estimated using the abbreviated Modification of Diet in Renal Disease (MDRD) equation,⁴⁹ and the results of the analysis were the same in terms of relative importance of predictors in determining outcome. Start of follow-up was defined as the first assessment suggestive of renal disease. A CrCl <15 ml/min per 1.73 m², initiation of dialysis, or transplantation defined ESRD. MAP was defined as the diastolic pressure (in mmHg) plus one third of the pulse pressure. For each patient, an average MAP was

determined for each 6-mo block during follow-up; the average of every 6-mo period's MAP is represented by the TA-MAP. Proteinuria was measured by 24-h urine protein collection. In a similar manner to MAP, the TA-proteinuria represents an average of the mean of every 6-mo period's proteinuria measurements.

The rate of renal function decline is expressed as the slope of CrCl, which was obtained by fitting a straight line through the calculated CrCl using linear regression and the principal of least squares. This was plotted and visually examined for each patient. Periods of reversible acute renal failure (a rapid reduction and recovery in CrCl of $\geq 40\%$ within a 1-mo time frame) were removed from the calculation.

Statistical Analyses

Data were analyzed using Microsoft Excel (Redmond, WA) and SPSS software (SPSS, Chicago, IL). Normally distributed variables are expressed as means \pm SD and compared using *t* test or ANOVA as required. Nonparametric variables are expressed as median and range and compared using either Mann-Whitney *U* or Kruskal-Wallis test. Categorical variables were compared using a χ^2 test. All *P* values were two-tailed; *P* < 0.05 was considered statistically significant. Univariate followed by multivariate linear regression was used to determine independent predictors of slope. Clinically relevant parameters or variables significantly associated with slope by univariate analysis were included in the multivariate models. Because proteinuria distribution was skewed (at presentation and time averaged), log-transformed values were used in the regression analysis, and similar results in terms of the significance of proteinuria were obtained with non-transformed data (data not shown). Multivariate regression models were assessed by stepwise and block entry of variables. Renal survival times were calculated from the first clinical assessment suggestive of renal disease to last follow-up. The relationship between parameters and renal survival was assessed using Cox regression.

Exposure to ACEi or ARB was considered as both a dichotomous variable and a continuous TA measurement of ACEi/ARB exposure. This variable had a skewed distribution and was considered in quartiles of exposure to ACEi/ARB for multivariate regression. For assessment of the role of ACEi/ARB on survival, these factors were considered as time-dependent variables for Cox regression analysis to account for time of initiation as well as residual GFR at that moment.

ACKNOWLEDGMENTS

H.N.R. is the recipient of a KRESCENT clinician-scientist fellowship from the Kidney Foundation of Canada and the Canadian Institute of Health Research (CIHR). J.W.S. holds the Amgen-CIHR Research Chair at University Health Network/University of Toronto. Funding for this study was supported by operating grants from the Kidney Foundation of Canada and CIHR, in addition to the CIHR Genes, Gender, and Glomerulonephritis New Emerging Team grant. S.T.'s research efforts are supported by the Fonds de la recherche en santé du Québec.

We thank the glomerulonephritis registrars N. Ryan and P. Ling for help in the collection and management of data and the following nephrologists for contributing patients and support to the registry:

Drs. S. Albert, J. Bargman, M. Berall, W. Berry, H. Bornstein, G. Buldo, C.J. Cardella, C. Chan, P. Chan, S. Chow, E.H. Cole, S. Donnelly, I.O. Elkan, S.S.A. Fenton, M.B. Goldstein, R. Golush, G. Hercz, M. Hladunewich, M.R. Hockley, V. Jassal, K. Kamel, A. Kang, S.Y. Karanickolas, D. Kim, L. Lam, A.G. Logan, C.E. Lok, M.E. Manuel, P. McFarlane, H. Mehta, D. Mendelssohn, D. Naimark, B. Nathoo, P.S.Y. Ng, M. Oliver, D.G. Oreopoulos, S. Pandeya, Y. Pei, Y.A. Pierattos, V. Pouloupoulos, R. Prasad, B. Reen, R.M. Richardson, J. Roscoe, D. Ryan, J. Sachdeva, C.S. Saiphoo, D. Sapir, J. Sasal, M. Schreiber, M. Silverman, A. Steele, E. Szaky, P. Tam, D.S. Thompson, R. Ting, S. Tobe, A. Wadgymar, L. Warner, C. Wei, C. Whiteside, G. Wong, G. Wu, and J. Zaltzman. We also thank participating pathologists T. Feltis, A.M. Herzenberg, S. Jothy, G. Lajoie, L. Sugar, and J. Sweet.

DISCLOSURES

None.

REFERENCES

- Maisonneuve P, Agodoa L, Gellert R, Stewart JH, Bucciati G, Lowenfels AB, Wolfe RA, Jones E, Disney AP, Briggs D, McCredie M, Boyle P: Distribution of primary renal diseases leading to end-stage renal failure in the United States, Europe, and Australia/New Zealand: Results from an international comparative study. *Am J Kidney Dis* 35: 157–165, 2000
- Bartosik LP, Lajoie G, Sugar L, Cattran DC: Predicting progression in IgA nephropathy. *Am J Kidney Dis* 38: 728–735, 2001
- D'Amico G, Minetti L, Ponticelli C, Fellin G, Ferrario F, Barbiano DB, Imbasciati E, Ragni A, Bertoli S, Fogazzi G: Prognostic indicators in idiopathic IgA mesangial nephropathy. *Q J Med* 59: 363–378, 1986
- Koyama A, Igarashi M, Kobayashi M: Natural history and risk factors for immunoglobulin A nephropathy in Japan. Research Group on Progressive Renal Diseases. *Am J Kidney Dis* 29: 526–532, 1997
- Radford MG Jr, Donadio JV Jr, Bergstralh EJ, Grande JP: Predicting renal outcome in IgA nephropathy. *J Am Soc Nephrol* 8: 199–207, 1997
- D'Amico G: Natural history of idiopathic IgA nephropathy: Role of clinical and histological prognostic factors. *Am J Kidney Dis* 36: 227–237, 2000
- Geddes CC, Rauta V, Gronhagen-Riska C, Bartosik LP, Jardine AG, Ibels LS, Pei Y, Cattran DC: A tricontinental view of IgA nephropathy. *Nephrol Dial Transplant* 18: 1541–1548, 2003
- D'Amico G: The commonest glomerulonephritis in the world: IgA nephropathy. *Q J Med* 64: 709–727, 1987
- Ibels LS, Gyory AZ: IgA nephropathy: Analysis of the natural history, important factors in the progression of renal disease, and a review of the literature. *Medicine (Baltimore)* 73: 79–102, 1994
- Szeto CC, Lai FM, To KF, Wong TY, Chow KM, Choi PC, Lui SF, Li PK: The natural history of immunoglobulin A nephropathy among patients with hematuria and minimal proteinuria. *Am J Med* 110: 434–437, 2001
- Usui J, Yamagata K, Kai H, Outeki T, Yamamoto S, Muro K, Hirayama A, Yoh K, Tomida C, Hirayama K, Suzuki S, Kobayashi M, Nagata M, Koyama A: Heterogeneity of prognosis in adult IgA nephropathy, especially with mild proteinuria or mild histological features. *Intern Med* 40: 697–702, 2001
- Li PK, Ho KK, Szeto CC, Yu L, Lai FM: Prognostic indicators of IgA nephropathy in the Chinese: Clinical and pathological perspectives. *Nephrol Dial Transplant* 17: 64–69, 2002
- Rauta V, Finne P, Fagerudd J, Rosenlof K, Tornroth T, Gronhagen-

- Riska C: Factors associated with progression of IgA nephropathy are related to renal function: A model for estimating risk of progression in mild disease. *Clin Nephrol* 58: 85–94, 2002
14. D'Amico G: Natural history of idiopathic IgA nephropathy and factors predictive of disease outcome. *Semin Nephrol* 24: 179–196, 2004
 15. Lee HS, Lee MS, Lee SM, Lee SY, Lee ES, Lee EY, Park SY, Han JS, Kim S, Lee JS: Histological grading of IgA nephropathy predicting renal outcome: Revisiting H. S. Lee's glomerular grading system. *Nephrol Dial Transplant* 20: 342–348, 2005
 16. Ruggenenti P, Perna A, Mosconi L, Pisoni R, Remuzzi G: Urinary protein excretion rate is the best independent predictor of ESRF in non-diabetic proteinuric chronic nephropathies. "Gruppo Italiano di Studi Epidemiologici in Nefrologia" (GISEN). *Kidney Int* 53: 1209–1216, 1998
 17. Donadio JV, Bergstralh EJ, Grande JP, Rademcher DM: Proteinuria patterns and their association with subsequent end-stage renal disease in IgA nephropathy. *Nephrol Dial Transplant* 17: 1197–1203, 2002
 18. Remuzzi G, Bertani T: Is glomerulosclerosis a consequence of altered glomerular permeability to macromolecules? *Kidney Int* 38: 384–394, 1990
 19. Zoja C, Donadelli R, Colleoni S, Figliuzzi M, Bonazzola S, Morigi M, Remuzzi G: Protein overload stimulates RANTES production by proximal tubular cells depending on NF-kappa B activation. *Kidney Int* 53: 1608–1615, 1998
 20. Morigi M, Macconi D, Zoja C, Donadelli R, Buelli S, Zanchi C, Ghilardi M, Remuzzi G: Protein overload-induced NF-kappaB activation in proximal tubular cells requires H(2)O(2) through a PKC-dependent pathway. *J Am Soc Nephrol* 13: 1179–1189, 2002
 21. Zoja C, Benigni A, Remuzzi G: Cellular responses to protein overload: Key event in renal disease progression. *Curr Opin Nephrol Hypertens* 13: 31–37, 2004
 22. Tang S, Leung JC, Abe K, Chan KW, Chan LY, Chan TM, Lai KN: Albumin stimulates interleukin-8 expression in proximal tubular epithelial cells in vitro and in vivo. *J Clin Invest* 111: 515–527, 2003
 23. Reich H, Tritschler D, Herzenberg AM, Kassiri Z, Zhou X, Gao W, Scholey JW: Albumin activates ERK via EGF receptor in human renal epithelial cells. *J Am Soc Nephrol* 16: 1266–1278, 2005
 24. Jafar TH, Stark PC, Schmid CH, Landa M, Maschio G, de Jong PE, de Zeeuw D, Shahinfar S, Toto R, Levey AS: Progression of chronic kidney disease: The role of blood pressure control, proteinuria, and angiotensin-converting enzyme inhibition—A patient-level meta-analysis. *Ann Intern Med* 139: 244–252, 2003
 25. Eiro M, Katoh T, Kuriki M, Asano K, Watanabe K, Watanabe T: The product of duration and amount of proteinuria (proteinuria index) is a possible marker for glomerular and tubulointerstitial damage in IgA nephropathy. *Nephron* 90: 432–441, 2002
 26. Troyanov S, Wall CA, Miller JA, Scholey JW, Cattran DC: Idiopathic membranous nephropathy: Definition and relevance of a partial remission. *Kidney Int* 66: 1199–1205, 2004
 27. Troyanov S, Wall CA, Miller JA, Scholey JW, Cattran DC: Focal and segmental glomerulosclerosis: Definition and relevance of a partial remission. *J Am Soc Nephrol* 16: 1061–1068, 2005
 28. Kobayashi Y, Tateno S, Hiki Y, Shigematsu H: IgA nephropathy: Prognostic significance of proteinuria and histological alterations. *Nephron* 34: 146–153, 1983
 29. Frimat L, Briancon S, Hestin D, Aymard B, Renoult E, Huu TC, Kessler M: IgA nephropathy: Prognostic classification of end-stage renal failure. L'Association des Nephrologues de l'Est. *Nephrol Dial Transplant* 12: 2569–2575, 1997
 30. Remuzzi A, Peticucci E, Ruggenenti P, Mosconi L, Limonta M, Remuzzi G: Angiotensin converting enzyme inhibition improves glomerular size-selectivity in IgA nephropathy. *Kidney Int* 39: 1267–1273, 1991
 31. Rekola S, Bergstrand A, Bucht H: Deterioration rate in hypertensive IgA nephropathy: Comparison of a converting enzyme inhibitor and beta-blocking agents. *Nephron* 59: 57–60, 1991
 32. Cattran DC, Greenwood C, Ritchie S: Long-term benefits of angiotensin-converting enzyme inhibitor therapy in patients with severe immunoglobulin a nephropathy: A comparison to patients receiving treatment with other antihypertensive agents and to patients receiving no therapy. *Am J Kidney Dis* 23: 247–254, 1994
 33. Maschio G, Cagnoli L, Claroni F, Fusaroli M, Ruggiu C, Sanna G, Sasdelli M, Zuccala A, Zucchelli P: ACE inhibition reduces proteinuria in normotensive patients with IgA nephropathy: A multicentre, randomized, placebo-controlled study. *Nephrol Dial Transplant* 9: 265–269, 1994
 34. Randomised placebo-controlled trial of effect of ramipril on decline in glomerular filtration rate and risk of terminal renal failure in proteinuric, non-diabetic nephropathy. The GISEN Group (Gruppo Italiano di Studi Epidemiologici in Nefrologia). *Lancet* 349: 1857–1863, 1997
 35. Maschio G, Alberti D, Janin G, Locatelli F, Mann JF, Motolese M, Ponticelli C, Ritz E, Zucchelli P: Effect of the angiotensin-converting-enzyme inhibitor benazepril on the progression of chronic renal insufficiency. The Angiotensin-Converting-Enzyme Inhibition in Progressive Renal Insufficiency Study Group. *N Engl J Med* 334: 939–945, 1996
 36. Russo D, Pisani A, Balletta MM, De Nicola L, Savino FA, Andreucci M, Minutolo R: Additive antiproteinuric effect of converting enzyme inhibitor and losartan in normotensive patients with IgA nephropathy. *Am J Kidney Dis* 33: 851–856, 1999
 37. Nakao N, Yoshimura A, Morita H, Takada M, Kayano T, Ideura T: Combination treatment of angiotensin-II receptor blocker and angiotensin-converting-enzyme inhibitor in non-diabetic renal disease (COOPERATE): A randomised controlled trial. *Lancet* 361: 117–124, 2003
 38. Praga M, Gutierrez E, Gonzalez E, Morales E, Hernandez E: Treatment of IgA nephropathy with ACE inhibitors: A randomized and controlled trial. *J Am Soc Nephrol* 14: 1578–1583, 2003
 39. Li PK, Leung CB, Chow KM, Cheng YL, Fung SK, Mak SK, Tang AW, Wong TY, Yung CY, Yung JC, Yu AW, Szeto CC: Hong Kong study using valsartan in IgA nephropathy (HKVIN): A double-blind, randomized, placebo-controlled study. *Am J Kidney Dis* 47: 751–760, 2006
 40. Remuzzi A, Perico N, Sangalli F, Vendramin G, Morigi M, Ruggenenti P, Remuzzi G: ACE inhibition and ANG II receptor blockade improve glomerular size-selectivity in IgA nephropathy. *Am J Physiol* 276: F457–F466, 1999
 41. Ruggenenti P, Perna A, Gherardi G, Garini G, Zoccali C, Salvadori M, Scolari F, Schena FP, Remuzzi G: Renoprotective properties of ACE-inhibition in non-diabetic nephropathies with non-nephrotic proteinuria. *Lancet* 354: 359–364, 1999
 42. Jafar TH, Schmid CH, Landa M, Giatras I, Toto R, Remuzzi G, Maschio G, Brenner BM, Kamper A, Zucchelli P, Becker G, Himmelmann A, Bannister K, Landais P, Shahinfar S, de Jong PE, de Zeeuw D, Lau J, Levey AS: Angiotensin-converting enzyme inhibitors and progression of nondiabetic renal disease: A meta-analysis of patient-level data. *Ann Intern Med* 135: 73–87, 2001
 43. Woo KT, Lau YK, Wong KS, Chiang GS: ACEI/ATRA therapy decreases proteinuria by improving glomerular permselectivity in IgA nephritis. *Kidney Int* 58: 2485–2491, 2000
 44. Haas M: Histologic subclassification of IgA nephropathy: A clinicopathologic study of 244 cases. *Am J Kidney Dis* 29: 829–842, 1997
 45. Bogenschutz O, Bohle A, Batz C, Wehrmann M, Pressler H, Kendziorra H, Gartner HV: IgA nephritis: On the importance of morphological and clinical parameters in the long-term prognosis of 239 patients. *Am J Nephrol* 10: 137–147, 1990
 46. Regional program for the study of glomerulonephritis. Central Committee of the Toronto Glomerulonephritis Registry. *Can Med Assoc J* 124: 158–161, 1981
 47. Cockcroft DW, Gault MH: Prediction of creatinine clearance from serum creatinine. *Nephron* 16: 31–41, 1976
 48. K/DOQI clinical practice guidelines for chronic kidney disease: Evaluation, classification, and stratification. *Am J Kidney Dis* 39: S1–S266, 2002
 49. Levey AS, Greene T, Kusek JW, Beck GJ: A simplified equation to predict glomerular filtration rate from serum creatinine [Abstract]. *J Am Soc Nephrol* 11: A0828, 2000